's	Seizure	Action	Plan
· ·	DCIZUIC	1 LC LI OII	1 1411

Child's Name

Date of Birth:					
Parent: Pl		none:			
Physician:	Phone:				
Physician Signature:	Date:				
<b>Seizure Information</b>					
Seizure Type	Length	Frequ	uency	Description	
Seizure triggers or warning signs:  Response after seizure:					
Special Considerations and Precautions: (activities, trips, diet)					
Treatment					
☐ Absence ☐ Atonic ☐ Complex Partial ☐ Infantile Spasms	<ol> <li>Stay with the child during and after the seizure. Although the child may appear conscious, he/she may lose awareness of surroundings.</li> <li>Be prepared to assist child to the floor if he loses consciousness.</li> <li>Document seizure in log.</li> <li>Notify parent.</li> <li>Special Instructions:</li> </ol>				
☐ General Tonic/Clonic	<ol> <li>Do not restrain movement. Let the seizure run its course.</li> <li>Turn child on side. Loosen the child's collar.</li> <li>Do not place anything in the mouth. Remove hard, sharp objects from the area.</li> <li>If possible turn head to the side in the event he/she vomits. (Use "Universal Precautions" if child vomits.)</li> <li>Observe, note time &amp; be prepared to describe the pattern of the seizure.</li> <li>Record details as they occur, or as soon as possible thereafter.</li> <li>Notify parent.</li> <li>When seizure is over, allow the child to rest.</li> <li>Stay with the child until fully recovered or parent arrives.</li> </ol>				
Administer Emergency Medication:	Diastat order:				
Emergency Response  Call 911!	Call 911 if:  the seizure lasts more than minutes, or  the child has a continuous seizure, or  the child remains unconscious after the seizure, or  he or she is having difficulty breathing, or  any injury resulted from the seizure.				

